www.chemistryhk.com

Write your name here Surname		Other names	5
Pearson Edexcel International Advanced Level	Centre Number		Candidate Number
Chamictus	_		
Chemistry Advanced Subsidiar Unit 2: Application of	ry	ciples	of Chemistry
Advanced Subsidiar	ry of Core Prin	ciples	of Chemistry Paper Reference WCH02/01

Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.

Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.
- Questions labelled with an asterisk (*) are ones where the quality of your written communication will be assessed
 - you should take particular care on these questions with your spelling, punctuation and grammar, as well as the clarity of expression, on these questions.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- Show all your working in calculations and include units where appropriate.

Turn over ▶







DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

SECTION A

Answer ALL the questions in this section. You should aim to spend no more than 20 minutes on this section. For each question, select one answer from A to D and put a cross in the box ⋈. If you change your mind, put a line through the box ⋈ and then mark your new answer with a cross ⋈.

- 1 Which compound is likely to show the **most** ionic character?
 - PH₃
 - B BH₃
 - C NaI
 - ☑ D KCl

(Total for Question 1 = 1 mark)

- **2** Which species has a similar shape to that of an ammonia molecule?
 - A BH₃
 - B CH₃⁺

 - \square **D** CO_3^{2-}

(Total for Question 2 = 1 mark)

- **3** Which compound contains a dative covalent bond?
 - A NH₃
 - B NCl₃

 - D CH₃NH₂


(Total for Question 3 = 1 mark)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

4 The diagram shows a liquid leaving a burette and passing a charged rod.



Which of the following liquids will be **most** attracted to the charged rod?

- A CCl₄
- B C₅H₁₂
- C CS₂
- ☑ D CHCl₃

(Total for Question 4 = 1 mark)

- **5** Which substance has more than one type of intermolecular force between its molecules in the liquid state?
 - A Br₂
 - \square **B** P_4
 - C NH₃
 - ☑ D CH₄

(Total for Question 5 = 1 mark)

DO NOT WRITE IN THIS AREA

	\boxtimes A	A	Graphite
	×	3	Diamond
	X (-	Cyclohexane
	X)	Buckminsterfullerene
			(Total for Question 6 = 1 mark)
7	Whic	:h	species contains bond angles equal to 90°?
			BeCl ₂
	× E		NH_4^+
	X (_	SiCl ₄
	× c)	SF_6
			(Total for Question 7 = 1 mark)
8			e solid gives a lilac flame-test colour. It reacts with water, forming a ly alkaline solution.
8	stror	ng	· · · · · · · · · · · · · · · · · · ·
8	stror The	ng so	ly alkaline solution.
8	stror The	ng so 1	ly alkaline solution.
8	stror	ng so A	ly alkaline solution. lid could be calcium oxide
8	The s	ng so A	ly alkaline solution. lid could be calcium oxide potassium oxide

4



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

	⊠ A	first ionisation energy decreases.
	В	atomic radius decreases.
	⊠ C	electronegativity increases.
	X D	number of outer shell electrons increases.
		(Total for Question 9 = 1 mark)
10 V	Vhich	Group 2 hydroxide is the most soluble in water?
	X A	Barium hydroxide
[В	Calcium hydroxide
	⊠ C	Magnesium hydroxide
	⊠ D	Strontium hydroxide
		(Total for Question 10 = 1 mark)
 []	× A	s the trend in bond energies for the sequence of molecules chlorine to bromine to iodine. Decreases Decreases to bromine then increases
[⊠ C	Increases
	⊠ D	Increases to bromine then decreases
		(Total for Question 11 = 1 mark)
		s seen when concentrated sulfuric acid is added to solid sodium chloride at emperature?
[X A	Green gas
	⊠ B	Steamy fumes
	⊠ C	White smoke
	⊠ D	Yellow solid




DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

13 Curve **X** was obtained when 0.50 g of calcium carbonate **powder** reacted with excess dilute hydrochloric acid at 20 °C.

Which curve best represents the reaction of a single 0.25 g **chip** of calcium carbonate with excess of the same dilute hydrochloric acid at the same temperature?



- A
- \boxtimes B
- X C
- X D

(Total for Question 13 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.


6

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

14 Consider the following reaction profile.



Which energy change would alter if a catalyst were added to the reaction?

- A
- \bowtie B
- X C
- **⋈** D


(Total for Question 14 = 1 mark)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

15 The Maxwell–Boltzmann distribution for a reaction mixture is shown below. E_a is the activation energy and N is the number of molecules with the most probable energy.



What is the effect of **increasing** the temperature on E_a and on N?

		Effect on E _a	Effect on N
X	A	increases	decreases
X	В	increases	increases
X	C	constant	decreases
X	D	constant	increases

(Total for Question 15 = 1 mark)

16 Which statement regarding a chemical reaction at equilibrium is **always** true?

- ☑ A The rates of the forward and backward reactions are equal.
- **B** The concentrations of reactants and products are equal.
- ☑ C The forward and backward reactions have stopped.
- **D** The addition of a catalyst changes the position of equilibrium.

(Total for Question 16 = 1 mark)

NOT WRITE IN THIS AREA

WRITE IN THIS AREA

- 17 In which system will a change in pressure have **no** effect on the position of equilibrium?
 - \square **A** $2NO_2(g) \rightleftharpoons N_2O_4(g)$
 - \square **B** $H_2(q) + I_2(q) \rightleftharpoons 2HI(q)$
 - \square **C** 2NO(g) + O₂(g) \rightleftharpoons 2NO₂(g)
 - \square **D** $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$

(Total for Question 17 = 1 mark)

- **18** 1.00 dm³ of a solution with a **sodium** ion concentration of 0.100 mol dm⁻³ is required. What volume of 0.200 mol dm⁻³ sodium sulfate solution is needed to make this solution by dilution with water?
 - \triangle **A** 100 cm³
 - \square **B** 250 cm³
 - \boxtimes **C** 500 cm³
 - \square **D** 1000 cm³

(Total for Question 18 = 1 mark)

19 Propan-2-ol is produced from the reaction of propene and steam. Assuming a 65% yield, what is the mass of propene required to produce 390 g of propan-2-ol?

[Molar masses / g mol⁻¹ propene = 42.0 propan-2-ol = 60.0]

- **B** 273 g

(Total for Question 19 = 1 mark)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

20	Pro	pei	ne can be formed by heating 1-bromopropane with alcoholic potassium hydroxide solution.
	Thi	is re	action is an example of
	X	A	reduction.
	×	В	hydrolysis.
	X	C	elimination.

condensation.

(Total for Question 20 = 1 mark)

TOTAL FOR SECTION A = 20 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

SECTION B

Answer ALL the questions. Write your answers in the spaces provided.

- 21 There are **four** isomeric alcohols with the molecular formula $C_4H_{10}O$.
 - (a) Two of these alcohols are butan-1-ol ($CH_3CH_2CH_2CH_2OH$) and butan-2-ol. The other two isomers are alcohol **X** and alcohol **Y**.
 - (i) Draw the **displayed** formula for butan-2-ol, showing all the bonds.

(1)

(ii) Alcohol **X** does not react with acidified potassium dichromate(VI) solution. Give the structure of alcohol **X**.

(1)


(iii) Name the fourth isomer, alcohol Y.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(b) The infrared spectrum of one of the four alcohols with the formula $C_4H_{10}O$ is shown.



Some infrared data is given in the table below.

Bond	Wavenumber / cm ⁻¹
C—H stretch, alkane	2962 — 2853
C—H stretch, alkene	3100 — 3010
O—H stretch (weak), carboxylic acids	3300 — 2500
O—H stretch (broad), alcohols	3750 — 3200

(i) Circle the relevant part of the infrared spectrum which confirms that this isomer is an alcohol.

(1)

(ii) State how the infrared spectrum can be used to identify which one of the four alcohols with the formula $C_4H_{10}O$ is present.

(1)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

www.chemistryhk.com

 (ii) Draw the displayed formula of the organic product formed when butan-1-ol is heated under reflux with acidified potassium dichromate(VI) solution. (1) (iii) Draw the displayed formula of the organic product formed when butan-1-ol is gently heated with acidified potassium dichromate(VI) solution and the product distilled off as it is formed. 		(Total for Question 21 = 9 ma	arks)
 (ii) Draw the displayed formula of the organic product formed when butan-1-ol is heated under reflux with acidified potassium dichromate(VI) solution. (1) (iii) Draw the displayed formula of the organic product formed when butan-1-ol is gently heated with acidified potassium dichromate(VI) solution and the product distilled off as it is formed. 	Гуре с		(1)
 (ii) Draw the displayed formula of the organic product formed when butan-1-ol is heated under reflux with acidified potassium dichromate(VI) solution. (1) (iii) Draw the displayed formula of the organic product formed when butan-1-ol is gently heated with acidified potassium dichromate(VI) solution and the 		(iv) State the type of reaction butan-1-ol undergoes in both (c)(ii) and (c)(iii).	
(ii) Draw the displayed formula of the organic product formed when butan-1-ol is heated under reflux with acidified potassium dichromate(VI) solution.		is gently heated with acidified potassium dichromate(VI) solution and the	(1)
(1)		• • • • • • • • • • • • • • • • • • • •	(1)
(i) Classify the alcohol butan-1-ol.		(i) Classify the alcohol butan-1-ol.	(1)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

- 22 This question is about halogenoalkanes.
 - (a) A student investigates the relative rate of hydrolysis of three halogenoalkanes.

The student mixes 5 cm³ of ethanol with five drops of halogenoalkane. This mixture is warmed to 50 °C in a water bath. The student adds 5 cm³ of aqueous silver nitrate, also heated to 50 °C, to the halogenoalkane. The time taken for a precipitate to form is recorded in a table.

Halogenoalkane	Time taken for a precipitate to form /s
CH ₃ CH ₂ CH ₂ Cl	120
CH ₃ CH ₂ CH ₂ CH ₂ Br	62
CH ₃ CH ₂ CH ₂ CH ₂ I	20

(i) Explain why ethanol is added to each halogenoalkane.

(1)

(ii) Complete the equation for the hydrolysis of 1-iodobutane, CH₃CH₂CH₂CH₂I.

(1)

$$CH_3CH_2CH_2CH_2I + H_2O \rightarrow$$

(iii) Give the **name** of the organic product which forms in each of these hydrolysis reactions.

(1)

(iv) Classify the type and mechanism of the reaction occurring when halogenoalkanes undergo hydrolysis, under the conditions described in this investigation.

(2)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

*(v) Describe and explain the trend in the rates of hydrolysis of the three halogenoalkanes.	(3)
	(3)
) Chlorofluorocarbons (CFCs) are halogenoalkanes which have been shown to play a role in ozone depletion. Halogenoalkanes, which contain fluorine as the only	
halogen (HFCs), do not harm the ozone layer.	
Following the Montreal Protocol of September 1987, CFCs have been replaced in	
many applications by HFCs.	
*(i) Explain why CFCs deplete the ozone layer, whereas HFCs do not.	
	(2)
(ii) Suggest a reason why ozone depletion still occurs.	
	(1)
(Total for Question 22 = 11 ma	rks)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

23 A student was asked to find the identity of a Group 1 metal hydroxide by titration.

The student was given the following outline method.

- Fill a burette with dilute hydrochloric acid
- Accurately weigh about 1.14 g of the metal hydroxide
- Carefully dissolve all the metal hydroxide in water, transfer the solution and washings to a volumetric flask and then add more water to make 250.0 cm³ of solution
- Thoroughly mix the contents of the volumetric flask
- Accurately transfer 25.0 cm³ of this solution to a conical flask
- Add two or three drops of a suitable indicator to the solution in the flask
- Carry out a rough titration of the solution in the flask with the dilute hydrochloric acid
- Accurately repeat the titration several times and calculate a mean titre.

The student's results and relevant data are shown.

Mass of metal hydroxide = 1.14q

Concentration of the hydrochloric acid = 0.730 g in 100 cm³ of solution

Mean titre = $23.80 \, \text{cm}^3$

(a) Give a reason why the student does not simply add 1.14 g of the metal hydroxide to 250 cm³ of water.

(1)

(b) Name the most suitable piece of apparatus for transferring 25.0 cm³ of the metal hydroxide solution to the conical flask.

(1)

(c) Name a suitable indicator to add to the solution in the conical flask and give the colour change at the end-point.

(3)

Colour change from ______ to _____



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(d) The equation for the reaction between the metal hydroxide and hydrochloric acid is shown. The letter **M** represents the unknown Group 1 metal.

$$MOH + HCl \rightarrow MCl + H_2O$$

(i) Calculate the concentration, in mol dm⁻³, of the hydrochloric acid in the burette.

(2)

(ii) Calculate the number of moles of hydrochloric acid used in the titration.

(1)

(iii) Deduce the number of moles of **M**OH in 25.0 cm³ of solution.

(1)

(iv) Calculate the total number of moles of **M**OH in the original solution.

(1)

(v) Calculate the molar mass of **M**OH, in g mol⁻¹.

(1)

(vi) Deduce the Group 1 metal present in the compound **M**OH.


(1)

(Total for Question 23 = 12 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

24 A hydrogen peroxide molecule can be represented by the structure shown.



(a) Suggest a value for the H—O—O bond angle.

(1)

- (b) Hydrogen peroxide dissolves in water.
 - (i) State the strongest type of interaction that occurs between molecules of hydrogen peroxide and water.

(1)

(ii) Draw a diagram to show the interaction named in (b)(i) between one molecule of hydrogen peroxide and one molecule of water.

Show any relevant lone pairs and dipoles in your diagram.

(3)

*(c) Explain, in terms of electronegativity, why the boiling temperature of H_2S_2 is lower than that of H_2O_2 .

(2)

(Total for Question 24 = 7 marks)

TOTAL FOR SECTION B = 39 MARKS



SECTION C

Answer ALL the questions. Write your answers in the spaces provided.

25 This question is about some aspects of the chemistry of Group 7 elements.

The elements in this group can also be described as p-block elements.

The elements chlorine, bromine and iodine can all be extracted from sea water.

The bromine in sea water is present as bromide ions, Br⁻.

The extraction of bromine from sea water occurs in four main stages.

- Stage 1: Formation of bromine when gaseous chlorine molecules are bubbled into an aqueous solution of bromide ions
- Stage 2: Removal of bromine vapour
- Stage 3: Production of hydrobromic acid solution from bromine
- Stage 4: Oxidation of hydrobromic acid to bromine.
- (a) State why the halogens are described as p-block elements.

//	-01	1
	7	
٧.		

(b) Give the physical states of bromine and iodine at room temperature.

(2)

Bromine

lodine



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(c) (i) Give an **ionic** equation for the reaction occurring in Stage **1**. Include state symbols in your answer.

(2)

(ii) Describe what you would **see** when the reaction in (c)(i) is carried out.

(1)

(iii) State the role of the chlorine during the reaction in (c)(i). Justify your answer in terms of electron transfer.

(2)

*(d) If Stage **1** is carried out under conditions of high acidity, this prevents the disproportionation of chlorine molecules.

$$Cl_2(g) + H_2O(I) \rightleftharpoons H^+(aq) + Cl^-(aq) + HClO(aq)$$

Explain, in terms of the above equilibrium, why this is so.

(2)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

www.chemistryhk.com

- (e) In Stage **3**, bromine vapour is converted into hydrobromic acid by the reaction of bromine with a mixture of sulfur dioxide and water vapour.
 - (i) Give the half-equation for the formation of bromide ions from bromine molecules. State symbols are not required.

(1)

(ii) Give the half-equation for the reaction between sulfur dioxide and water molecules to form hydrogen ions and sulfate(VI) ions.

State symbols are not required.

(1)

(iii) Hence write the ionic equation for the reaction between bromine and sulfur dioxide, in the presence of water, to form hydrogen ions, bromide ions and sulfate(VI) ions. State symbols are not required.

(2)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(f) (i) Identify, by name or formula, a compound that can be used to distinguish between separate aqueous solutions of potassium bromide and potassium ioc	dide. (1)
(ii) State what would be observed when a solution of the compound in (f)(i) is added to each of the separate aqueous solutions of potassium bromide and potassium iodide.	(2)
Observation with potassium bromide:	
Observation with potassium iodide:	
(iii) Identify, by name, a further reagent that could be added to the mixtures resulting from the test in (f)(ii) to confirm the identity of the halide ions.	(2)
(iv) Compare the observations that would be made when the reagent identified in (f)(iii) is added to the mixtures formed in (f)(ii).	(2)
(Total for Question 25 = 21 ma	nrks)
TOTAL FOR SECTION C = 21 MA TOTAL FOR PAPER = 80 MA	

P 5 0 7 8 6 A 0 2 2 2 4

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



The Periodic Table of Elements

0 (8)	(18) 4.0 He hetium	20.2
7	(17)	19.0
9	(16)	16.0
2	(15)	14.0
4	(13) (14) (15) (16) (17)	12.0
m	(13)	10.8
	1.0 H hydrogen 1	
		<u></u>
	Key	relative atomic mas
2	;) Key	.0 relative atomic mass
2) (2) Key	9 9.0 relative atomic mas

										_	П	_							٦				
2	707	Ne	neon	10	6'68	Ar	argon 18	83.8	Ā	krypton 24	စ္ခင္	131.3	Xe	xenon	54	[222]	R	radon 86	8		ted		
(17)	19.0	L	fluorine	6	32.5	บ	chlorine 17	6.62	Br	bromine 25	cs	126.9	_	iodine	53	[210]	Αt	astatine os	6		been repor		
(16)	16.0	0	oxygen		32.1	S	sulfur 16	79.0	Se	selenium 24	24	127.6	<u>a</u>	tellurium	52	[506]	S S	polonium 84	04		116 have !	nticated	
(15)	14.0	z	nitrogen	7	31.0	۵.	phosphorus 15	74.9	As		33	121.8	Sb	antimony	51	209.0	Bi	bismuth 92	63		mbers 112.	but not fully authenticated	
(14)	_		_		28.1	Si	silicon 14	72.6	g	germanium	32	118.7	Sn	ţi	50	207.2	Ъ	lead 82	70		atomic nu	but not f	
(13)	10.8	മ	boron	5	27.0	¥	aluminium 13	I		gallium	51	114.8	드	indium	49	204.4	F	thallium 81	5		Elements with atomic numbers 112-116 have been reported		
•							(12)	65.4	Zu	zinc	3	112.4	ੲ	cadminm	48	200.6	Ŧ	mercury	20		Elen		
							(11)	63.5	D C	copper	67	107.9	Ag	silver	47	١,			[1777]	[7/7]	Rg	roentgenium	111
							(10)	58.7	ï	nickel	87	106.4	В	palladium	46	195.1	చ	platinum 78	[274]		Ds	darmstadtium	110
							(6)	58.9	ပ	cobalt	/7	102.9	윤	rhodium	45	192.2		iridium 77		[007]		meitnerium	
-							(8)	55.8	Ā	iron		101.1	Ru	ruthenium	44	190.2	ŏ	osmium 76	1777	[//7]		hassium	
							(2)	54.9	Wn		Ç2	[86]	ပ	technetium	43	186.2	Re	rhenium 75	[7,24]	[+07]	Bh	pohrium	107
	mass	lod		nmper			(9)	52.0	ъ	vanadium chromium m	74	95.9	¥	5	42	183.8	>	tungsten 74	17767	[007]	Sg	seaborgium	106
Key	relative atomic mass	atomic symbol	name	atomic (proton) number			(2)	50.9	>	vanadium	73	92.9	g	iobium	41	180.9	٦a	tantalum 72	[7,77]	[707]	음	dubnit	105
	relati	ato		atomic			(4)	47.9	i=	titanium	77		Zr	zirconium	40	178.5	Ŧ	hafnium 72	[72,41]	[107]	¥	nutherfordium	5
							(3)	45.0	Sc	scandium	71	88.9	>	/ttrium	39	138.9	La*	lanthanum 57	[22,27]	[777]		actinium	
(2)	0.6	Be	beryllium	4	24.3	Mg	magnesium 12	40.1	g	calcium	07	97.8	Sr	strontium	38	137.3	Ва	barium 56	12,67	[077]	Ra	radium	88
(1)	6.9	ב	lithium	3	23.0	Na	sodium 11	39.1	¥	potassium	13	85.5	8				క	caesium 55	[222]	[677]	፫	francium	87
	_	_	_	_	_	_		_	_	_	_	_	_		_	_	_		_	_	_	_	_

175	Lu lutetium	71	[257]	۲	lawrenciun	103
173	Yb vtterbium	70	[254]	ž	nobelium	102
169	T m	69	[526]	Þ₩	mendelevium	101
167	Er erbium	68	[253]	FB	fermium	100
165	4	67		E	einsteinium	66
163	Dy dysprosium	, 66	[251]	უ	californium	86
159	Tb erbium	65	[245]	æ	berkelium	97
157	Gd adolinium	64	[247]	Ę	curium	96
152	Eu n europium	63	[243]	Αm	americium	95
150	Sm samarium	62	[242]	Pu	plutonium	94
[147]	Pm promethium	61	[237]	å	neptunium	93
144	Nd neodymium	60	238		Š	92
141	Pr praseodymium	59	[231]	Pa	protactinium	91
140	Ce cerium			두	thorium	06

* Lanthanide series * Actinide series