www.chemistryhk.com

Write your name here Surname	Othe	er names
Pearson Edexcel International Advanced Level	Centre Number	Candidate Number
Chamistry)
Chemistry Advanced Unit 5: General Principle Metals and Orga (including synop	es of Chemistry Inic Nitrogen Ch	emistry
Advanced Unit 5: General Principle	es of Chemistry Inic Nitrogen Ch Otic assessment)	Paper Reference
Advanced Unit 5: General Principle Metals and Orga (including synop	es of Chemistry Inic Nitrogen Ch Otic assessment)	emistry

Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.

Information

- The total mark for this paper is 90.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.
- Questions labelled with an asterisk (*) are ones where the quality of your written communication will be assessed
 - you should take particular care with your spelling, punctuation and grammar, as well as the clarity of expression, on these questions.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- Show all your working in calculations and include units where appropriate.

Turn over ▶

P50789A
©2017 Pearson Education Ltd.
6/5/5/5/5/1/





DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

SECTION A

Answer ALL the questions in this section. You should aim to spend no more than 20 minutes on this section. For each question, select one answer from A to D and put a cross in the box \boxtimes . If you change your mind, put a line through the box \boxtimes and then mark your new answer with a cross \boxtimes .

- 1 Which species contains an element with the same oxidation number that sulfur has in NaHSO₄?
 - \square **A** $K_4Fe(CN)_6$
 - B NH₄VO₃

 - \square **D** $[CoCl_4]^{2-}$

(Total for Question 1 = 1 mark)

- 2 Which species contains a bond angle of 90°?
 - \triangle **A** [Pt(NH₃)₂Cl₂]
 - \mathbf{B} [CuCl₂]
 - \square **C** $[CrCl_4]^-$
 - ☑ D SiCl₄

(Total for Question 2 = 1 mark)

- **3** Which equation can only be described as a ligand exchange reaction?
 - **A** $[Zn(OH)_2](s) + 2OH^-(aq) \rightarrow [Zn(OH)_4]^{2-}(aq)$
 - **B** $[Fe(H_2O)_6]^{3+}(aq) + H_2O(I) \rightarrow [Fe(H_2O)_5(OH)]^{2+}(aq) + H_3O^{+}(aq)$
 - \square **C** $[Cu(H_2O)_6]^{2+}(aq) + 2NH_3(aq) \rightarrow Cu(H_2O)_4(OH)_2(s) + 2NH_4^+(aq)$
 - \square **D** $[Cr(H_2O)_6]^{3+}(aq) + 6NH_3(aq) \rightarrow [Cr(NH_3)_6]^{3+}(aq) + 6H_2O(I)$

(Total for Question 3 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.

DO NOT WRITE IN THIS AREA

WRITE IN THIS AREA

4 A metal ion, M⁺, disproportionates spontaneously:

$$2M^+(aq) \rightarrow M(s) + M^{2+}(aq)$$

The table below gives the standard electrode potentials of four systems.

System	E [⊕] / V
$Cu^{2+}(aq) + e^- \rightarrow Cu^+(aq)$	+0.15
$Cu^+(aq) + e^- \rightarrow Cu(s)$	+0.52
$Ag^{2+}(aq) + e^{-} \rightarrow Ag^{+}(aq)$	+1.98
$Ag^{+}(aq) + e^{-} \rightarrow Ag(s)$	+0.80

Use the data to predict which statement is true under standard conditions.

- A Both Cu⁺ and Ag⁺ can disproportionate.
- B Only Cu⁺ can disproportionate.
- ☑ C Only Ag⁺ can disproportionate.
- ☑ D Neither Cu⁺ nor Ag⁺ can disproportionate.

(Total for Question 4 = 1 mark)

In a reaction of dichromate(VI) ions, $Cr_2O_7^{2-}$, with a metal ion M^{2+} , the oxidation number of each Cr atom changes from +6 to +3.

Each dichromate(VI) ion reacts with three M²⁺ ions.

What is the oxidation number of M after the reaction?

(1)

- **B** +3
- **◯** C +4
- **D** +5

(Total for Question 5 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.

DO NOT WRITE IN THIS AREA

A ligand exchange reaction occurs when EDTA is added to a solution containing the ion $[Cu(H_2O)_2(NH_3)_4]^{2+}$.

What is the **best** explanation for this?

- A A complex forms with copper and EDTA which is more soluble than the original.
- **B** A complex forms with copper and EDTA which is less soluble than the original.
- \triangle $\triangle H_{\text{reaction}}$ is positive for the reaction of $[Cu(H_2O)_2(NH_3)_4]^{2+}$ with EDTA.
- \square **D** ΔS_{system} is positive for the reaction of $[\text{Cu}(\text{H}_2\text{O})_2(\text{NH}_3)_4]^{2+}$ with EDTA.

(Total for Question 6 = 1 mark)

7 The repeat units of four polymers are shown:

Polymer	Repeat unit
Р	— CH₂CH(CONH₂) —
Q	— HN(CH₂)₅NHOC(CH₂)₃CO —
R	— HNCH₂CONHCH(CH₃)CO —
S	— CH₂CH(CN) —

(a) Which formula shows the repeat unit of the polymer poly(propenamide)?

(1)

- A P
- \square **B** Q

- (b) Which formula shows the repeat unit of a polymer which could be made using a diamine and a dioic acid as the monomers?

(1)

- A P
- \boxtimes **B** Q
- D S

(Total for Question 7 = 2 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

8 The enthalpy changes for the hydrogenation of cyclohexene and of benzene are shown:

$$C_6H_{10} + H_2 \rightarrow C_6H_{12}$$
 $\Delta H = -120 \text{ kJ mol}^{-1}$

$$C_6H_6 + 3H_2 \rightarrow C_6H_{12}$$
 $\Delta H = -208 \text{ kJ mol}^{-1}$

The delocalised structure of benzene is more stable than a molecule with three C=C double bonds by

- B 328 kJ mol⁻¹
- D 88 kJ mol⁻¹

(Total for Question 8 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.

The formula of the zwitterion of serine is

(Total for Question 9 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.

DO NOT WRITE IN THIS AREA

10 This question is about the amino acids, glycine and alanine.

$$H_2N$$
— C — $COOH$ H_2N — C — $COOH$ H_3

glycine

alanine

What is the total number of different dipeptides which can be made by reacting each of the optical isomers (enantiomers) of alanine with glycine?

- **B** 3
- **区** 4
- □ B

(Total for Question 10 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

11 The structure of the asthma medication salbutamol is

(a) Which functional group is **not** present in salbutamol?

(1)

- A Alcohol
- B Amide
- C Amine
- ☑ D Phenol
- (b) What is the molecular formula of salbutamol?

(1)

- \square A $C_{13}H_{22}NO_3$
- \blacksquare **B** $C_{13}H_{21}NO_3$
- \square **C** $C_{13}H_{20}NO_3$
- \square **D** $C_{13}H_{19}NO_3$

(Total for Question 11 = 2 marks)

12 The compound **X** with molecular formula $C_7H_{14}O_2$ shown is responsible for some of the flavour of bananas.

compound X

(a) The number of peaks in the **low** resolution proton nmr spectrum of **X** is

(1)

- A 4
- **■ B** 5
- **C** 6
- D 14



DO NOT WRITE IN THIS AREA

(b) Th	e number of singlets in the high resolution proton nmr spectrum of X is	(1)
⊠ A	1	,
⊠ B	2	
⊠ C	3	
⊠ D	5	
in	sample of X is hydrolysed with dilute acid. Which of these peaks is present the infrared spectrum of X but is not present in the spectra of its hydrolysis oducts?	
Re	fer to the correlation table on page 6 of the Data Booklet.	
		(1)
⊠ A	2962–2853 cm ⁻¹	
⋈ B	1750–1735 cm ⁻¹	
	1725–1700 cm ⁻¹	
⊠ D	1485–1365 cm ⁻¹	
(d) In	the mass spectrum of \mathbf{X} , which m/e peak would you expect \mathbf{not} to be present?	(1)
⊠ A	29	
	43	
	87	
⊠ D	129	
	(Total for Question 12 = 4 ma	rks)
	ple of 0.25 mol of a cyclic hydrocarbon produced 66 g of carbon dioxide and of water on combustion. The hydrocarbon is	
⊠ A	cyclohexane.	
⊠ B	cyclohexene.	
⊠ C	benzene.	
⊠ D	ethylbenzene.	



Aspirin Molar mass 180 g mol⁻¹

(a) In the preparation, 2.00 g of 2-hydroxybenzoic acid produced 1.65 g of aspirin.

The percentage yield was

(1)

- 92.9
- 82.5
- 76.7
- 63.3
- (b) The aspirin is purified by recrystallisation using water as the solvent. Which of the following statements is true?

(1)

- ☑ A The aspirin should be dissolved in an excess of water.
- The solution should be heated to above the melting temperature of aspirin.
- Soluble impurities are removed by filtering the mixture after cooling.
- Only water soluble impurities in the aspirin can be removed.

(Total for Question 14 = 2 marks)

TOTAL FOR SECTION A = 20 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

SECTION B

Answer ALL the questions. Write your answers in the spaces provided.

15 This question is about successive elements in Period 4.

	Potassium	Calcium	Scandium	Titanium
Atomic radius / nm	0.235	0.197	0.164	0.147
Sum of first and second ionisation energies / kJ mol ⁻¹	3470	1735	1866	1968

*(a)	There is a decrease in atomic radius between potassium and calcium, and also
	between scandium and titanium. Explain why the atomic radius decreases in
	both cases, and why the decrease between potassium and calcium is greater.

(2)

*(b)	Explain why the sums of the first two ionisation energies of Ca, Sc and Ti are al
	similar but less than the sum of the first two ionisation energies of potassium.

(2)



but scandium is not a transition element.	Periodic Table,
	(2)
(d) Aqueous solutions of Ti(II) compounds contain the ion $[Ti(H_2O)_6]^{2+}$.	
Draw a diagram of this ion.	
Name its shape, state the bond angle between adjacent water molecules type of bond between the ligand and the titanium.	and the
type of bond between the figure and the titalian.	(2)
(Total for Question 1	5 = 8 marks)
(Total for Question 1	5 = 8 marks)

- **16** This question is about manganese and its ions.
 - (a) (i) Complete the electronic configuration of Mn and Mn²⁺.

(1)

Mn [Ar]

Mn²⁺ [Ar]

(ii) Suggest why the oxidation number +2 is very stable for manganese.

(1)

(b) Consider the data below.

Electrode system	Standard electrode potential (E^{\ominus}) / V
$Mn^{2+}(aq) + 2e^- \rightleftharpoons Mn(s)$	-1.19
$MnO_4^-(aq) + 8H^+(aq) + 5e^- \rightleftharpoons Mn^{2+}(aq) + 4H_2O(I)$	+1.51

(i) Draw a labelled diagram showing how to set up a cell with the two electrode systems in the table above, in order to measure $E_{\text{cell}}^{\ominus}$.

State the standard conditions required for the cell.

(4)

DO NOT WRITE IN THIS AREA

(ii) Calculate the value of $E_{\text{cell}}^{\leftrightarrow}$.	(1)
(iii) Write the equation for the overall reaction that occurs in this cell State symbols are not required.	1. (2)
(c) A student investigated the redox chemistry of manganese by adding to a solution containing vanadium(III) ions, V ³⁺ (aq). Deduce what proin this reaction.	
Justify your answer by using the data in (b), and by considering item the table of Standard Electrode Potentials on pages 14 and 15 in the	
You should calculate E_{cell}^{\oplus} for any feasible reactions.	(4)

www.chemistryhk.com

- (d) A solution containing $\mathrm{Mn}^{2+}(\mathrm{aq})$ reacts with aqueous sodium hydroxide to form a precipitate.
 - (i) Write the **ionic** equation for the formation of the precipitate. Include state symbols.

(1)

(ii) State the colour of the precipitate.

(1)

(iii) The precipitate darkens if it is left standing in air. Identify, by name or formula, the new manganese compound responsible for this darkening.

(1)


(Total for Question 16 = 16 marks)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

17 The structure of a detergent is shown. In simplified form, the symbol R can be used to represent the alkyl group.



The detergent can be synthesised from benzene by the following route:

(a) (i) Give the number of carbon atoms and the number of hydrogen atoms in the alkyl group R.

(1)

(ii) Draw the structural formula of the **compound** which could be used to react with benzene in Step **1** and identify the catalyst required for this reaction.

(2)

Compound:

Catalyst:

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(iii) Write a mechanism for the alkylation of benzene in Step 1, using the symbol R for the alkyl group. You should include an equation showing the formation of the electrophile.


(3)

(iv) Name the reagent which is used to sulfonate the benzene ring in Step 2.

(1)

(v) Complete the diagram to show the **displayed** formula of the —SO₃H group in the product formed in Step **2**.

(1)



DO NOT WRITE IN THIS AREA

(vi)	ldentify a suitable reagent for use in Step 3 .	(1)
*(b) (i)	The product of Step 1 , C_6H_5R , is not soluble in water. Explain this fact by ident the intermolecular forces involved and comparing their relative strengths.	ifying
		(2)
(ii)	The detergent, $RC_6H_4SO_3Na$, is soluble in water. Explain how the bonding in this compound allows it to dissolve in water.	
		(2)
	(Total for Question 17 = 13 m	arks)

DO NOT WRITE IN THIS AREA

(a)		th benzene and phenol can be brominated in substitution reactions under table conditions.	
	(i)	Give the equation for the reaction of benzene with bromine. Include state symbols.	(2)
	(ii)	Give the equation for the reaction of phenol with bromine water, showing the structure of the organic product. State symbols are not required.	(1)
	*(iii)	Explain the effect the –OH group has on the bromination of the benzene ring.	(2)



DO NOT WRITE IN THIS AREA

('n) (i)	Phenol	reacts	with	dilute	nitric	acid
١	U,	, ,	11	LUGIIOI	reacts	VVILII	ullute	HILLIC	aciu.

Give the formula of **one** organic product of this reaction and state the type and mechanism of the reaction which is occurring.

(2)

Formula:

Type and mechanism:

(ii) Phenylamine also reacts with dilute nitric acid but the reaction does not involve the benzene ring.

Give the formula of the organic product in this reaction and state the type of reaction which is occurring.

(2)

Formula:

Type of reaction:

(c) Write the equation for the reaction of phenylamine with ethanoyl chloride, giving the **displayed** formula of the organic product.

(2)



RITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(d) Phenylamine is the starting material for making certain dyes.

Write equations to show how a dye could be made in two steps, using phenylamine and phenol as the only organic compounds. State the conditions needed for the first step.

(3)

First step:

Conditions

Second step:

(Total for Question 18 = 14 marks)

TOTAL FOR SECTION B = 51 MARKS



DO NOT WRITE IN THIS AREA

SECTION C

Answer ALL the questions. Write your answers in the spaces provided.

19

The element chromium occurs in the Earth's crust at a concentration of about 1.02×10^2 ppm. The name of the element comes from the Greek word for colour, and compounds of the three main oxidation states (+2, +3 and +6) are brightly coloured.


The element and its compounds have many uses, such as the manufacture of important alloys, pigments for paint and inks, colouring glass and tanning leather.

Chromium is an essential mineral for physiological functions but, above a certain level, its compounds are toxic. The target in many states of the USA is for a concentration in drinking water below 0.10 parts per billion. The maximum permissible level of Cr(VI) allowed to be released into waterways is 50 parts per billion. Chromium concentrations in drinking water can be measured by a colorimetric method.

*(a)	Solutions containing Cr ²⁺ (aq) and Cr ³⁺ (aq) are coloured due to electron transitions
	between d orbitals.

Explain why Cr²⁺(aq) and Cr³⁺(aq) **differ** in colour.

(2)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

www.chemistryhk.com

*(b) (i)	The oxide CrO_3 contains chromium in the oxidation state +6. Suggest two reasons why CrO_3 is unlikely to contain Cr^{6+} ions.	(2)
(ii)	Predict, by writing an equation, how CrO_3 would react with water.	(1)
(iii)	Write the equation for the reaction of dichromate(VI) ions with alkali in which chromate(VI) ions are formed. State symbols are not required.	(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

- (c) The concentration of ethanol in white wine can be measured using potassium dichromate(VI) titrations. A suitable method is:
- Step 1 A standard solution of ammonium iron(II) sulfate was used to measure the concentration of a potassium dichromate(VI) solution. The reactions are

$$Fe^{2+}(aq) \to Fe^{3+}(aq) + e^{-}$$

$$Cr_2O_7^{2-}(aq) + 14H^{+}(aq) + 6e^{-} \to 2Cr^{3+}(aq) + 7H_2O(I)$$

The concentration of the potassium dichromate(VI) solution was found to be $0.0210\,\mathrm{mol\,dm^{-3}}$.

Step 2 A mixture of 1.00 cm³ of wine, 100 cm³ of the potassium dichromate(VI) solution (an excess) and about 25 cm³ of dilute sulfuric acid were reacted until all the ethanol in the wine had been oxidised by the dichromate(VI) ions.

$$2Cr_2O_7^{2-}(aq) \,+\, 16H^+(aq) \,+\, 3C_2H_5OH(I) \,\rightarrow\, 4Cr^{3+}(aq) \,+\, 11H_2O(I) \,+\, 3CH_3COOH(aq)$$

- Step **3** The reaction mixture from Step **2** was made up to a volume of 200.0 cm³ and a portion was used to fill a burette. It was titrated with 25.00 cm³ portions of acidified ammonium iron(II) sulfate with concentration 0.015 mol dm⁻³. The mean titre was 18.60 cm³.
 - (i) An indicator of sodium diphenylamine and phosphoric acid was used to detect the end-point of the titrations.

If the titrations were carried out without an indicator, the colour change at the end-point would be difficult to detect. Explain, by reference to the ions involved and their colours, why this is the case.



(ii) Name the experimental procedure which would be used in the oxidation of ethanol to ethanoic acid in Step **2**.

(1)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(iii) How many moles of dichromate(VI) ions were present at the start of Step 2?

(iv) Use the results from Step **3** to calculate the number of moles of potassium dichromate(VI) solution present after it has oxidised the ethanol, and hence the number of moles used in the reaction.

(4)

(v) Calculate the number of moles of ethanol which were present in the $1.00\,\mathrm{cm^3}$ of wine.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(vi) The concentration of ethanol in wine is usually quoted as the percentage of "Alcohol by Volume" (ABV).

The volume of 1 mole of ethanol is 58.3 cm³.

Calculate the percentage of "Alcohol by Volume" in the wine.

(2)

26



WRITE IN THIS AREA

(d) The concentration of dichromate(VI) ions in drinking water can be measured by reacting them with a compound called diphenylcarbazide. This produces a complex with a reddish-purple colour. The concentration of this product is measured using a colorimeter.

The structure of diphenylcarbazide is shown.

The complex, which is thought to be octahedral, contains one chromium(III) ion formed by reduction.

Draw a circle round the atoms which are most likely to bond with the chromium ion. Hence predict the number of moles of diphenylcarbazide per mole of complex. Justify your answer.

(2)

(Total for Question 19 = 19 marks)

TOTAL FOR SECTION C = 19 MARKS TOTAL FOR PAPER = 90 MARKS



The Periodic Table of Elements

0 (8)

9

						٥											(18)
							1.0										4.0
							hvdrogen										¥.
	(2)			Key			. –					(13)	(14)	(12)	(16)	(17)	netium 2
\vdash	9.0	_	relati	relative atomic mass	mass							10.8	12.0	14.0	16.0	19.0	20.2
	Be		ato	atomic symbol	pol							8	U	z	0	L	N e
	beryllium		•	name								boron	carbon	nitrogen	oxygen	fluorine	neon
3	4		atomic	atomic (proton) number	number							5	9	7	8	6	10
ı	24.3										•	27.0	28.1	31.0	32.1	35.5	39.9
	W											¥	Si	۵	S	บ	Αr
sodium n	magnesium	(3)	9	(5)	(9)	6	(8)	(6)	(10)	(11)	(12)	aluminium 12	silicon	phosphorus	sulfur 16	chlorine 17	argon
+	40.1	45.0	47.9	50.9	52.0	54.9	55.8	58.9	58.7	63.5	65.4	69.7	72.6	74.9		79.9	83.8
¥	g	Sc	ï	>	ე	W	Fe	ပိ	ż	ŋ	Zn	Ga	g	As	Se	Br	궃
Ē	calcium	scandium	titanium	vanadium	romium	manganese	iron	cobalt	nickel	copper	zinc		germanium	arsenic	selenium	bromine	krypton
	20	21	22	23	24	25	76	27	28	59	30		32	33	34	35	36
	97.6	88.9	91.2	92.9	95.9	[86]	101.1	102.9	106.4	107.9	112.4	114.8	118.7	121.8	127.6	126.9	131.3
	Ş	>	Zr	S	Wo	ပ	Ru	뫈	Ь	Ag	5		Sn	Sb	<u>P</u>	Н	×
E	strontium	yttrium	zirconium	niobium	lybdenum	Ε	ruthenium	_	palladium	silver	cadmium		tin	antimony	tellurium	iodine	xenon
	38	39	4	41	42	43	4		46	47	48	49	20	51	52	53	24
132.9	137.3	138.9	178.5	180.9	183.8	186.2	190.2	192.2	195.1	197.0	200.6	204.4	207.2	209.0	[607]	[210]	[222]
	Ba	Ľa*	Ŧ	Б	>	Re	õ	Ļ	굽	Αn	Η	F	Ъ	Bi	&	Αt	~
	barium	lanthanum	hafnium	tantalum	tungsten	rhenium	osmium	iridium	platinum	plog	mercury	thallium	lead	bismuth	polonium	astatine	radon
\dashv	8	27	72	73	74	75	76	1	78	6/	80	81	82	83	84	85	86
[223]	[526]	[227] [261] [262]	[761]	[597]	[596]	[564]	[277]	[368]	[268] [271] [272]	[272]							
	Ra	Ac*	₹	<u>ප</u>	Sg	絽	Ұ	₩	Ds	Rg		ents with	atomic nu	mbers 112	-116 have t	Elements with atomic numbers 112-116 have been reported	pa:
۶	radiun	actinium	rutherfordium	dubnium	seaborgium	pohrium	hassium	meitnerium	damstadtium	roentgenium			but not f	but not fully authenticated	nticated		
_	88	68	104	105	106	107	108	109	110	111							

* Lanthanide series

175 **Lu** lutetium

173 **Yb** ytterbium

169 **Tm** thulium 69

167 **Er** erbium 68

165 **Ho** holmium

163

Dy
dysprosium
66

159 **Tb** terbium 65

152 **Eu** europium 63

samarinm

Pm romethium

141
Pr
aseodymium
59

[251]

Cf
alifornium
98

* Actinide series

Ce cerium 58 232 Th thorium 90