www.chemistryhk.com

Please check the examination details b	below before ente	ering your candidate information
Candidate surname		Other names
Pearson Edexcel International Advanced Level	entre Number	Candidate Number
Wednesday 9	Janua	ry 2019
Morning (Time: 1 hour 30 minutes)	Paper R	eference WCH01/01
Chemistry Advanced Subsidiary Unit 1: The Core Principle	es of Chen	nistry
Candidates must have: Scientific Ruler	calculator	Total Marks

Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.

Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.
- Questions labelled with an asterisk (*) are ones where the quality of your written communication will be assessed
 - you should take particular care with your spelling, punctuation and grammar, as well as the clarity of expression, on these questions.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ▶



P54457A ©2019 Pearson Education Ltd. 2/2/2/2/



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

SECTION A

Answer ALL the questions in this section. You should aim to spend no more than 20 minutes on this section. For each question, select one answer from A to D and put a cross in the box ⋈. If you change your mind, put a line through the box ⋈ and then mark your new answer with a cross ⋈.

- 1 The European Union limit for nitrogen dioxide in the atmosphere is 0.0209 cm³ in 1 m³. In units of parts per million (ppm) this is
 - \triangle **A** 2.09 × 10⁻⁵
 - **B** 2.09×10^{-2}
 - **C** 20.9
 - **D** 2.09×10^4

(Total for Question 1 = 1 mark)

- 2 A sample of blood plasma contains $3.10 \, \text{mg}$ of sodium ions in $1 \, \text{cm}^3$. The concentration, in mol dm⁻³, of sodium ions in the plasma is
 - \triangle **A** 1.35 × 10⁻¹
 - **B** 2.82×10^{-1}
 - \triangle **C** 1.35 × 10⁻⁴
 - \square **D** 2.82 × 10⁻⁴

(Total for Question 2 = 1 mark)

- **3** Dilute sulfuric acid is mixed with a solution of barium chloride. The reaction that occurs is

 - A displacement.B neutralisation.
 - C precipitation.
 - **D** redox.

(Total for Question 3 = 1 mark)



DO NOT WRITE IN THIS

DO NOT WRITE IN THIS

DO NOT WRITE IN THIS AREA

www.chemistryhk.com

4 How many **atoms** are there in 120 cm³ of ammonia gas at room temperature and pressure (r.t.p.)?

[Molar volume of gas at r.t.p. = $24\,000\,\mathrm{cm^3}\,\mathrm{mol^{-1}}$ Avogadro constant = $6.0\times10^{23}\,\mathrm{mol^{-1}}$]

- \triangle **A** 3.0 × 10²¹
- **B** 1.2×10^{22}
- \blacksquare **C** 1.5 × 10²²
- \square **D** 1.2 × 10²⁵

(Total for Question 4 = 1 mark)

5 The reaction of magnesium chloride with silver nitrate gives a precipitate of silver chloride.

$$MgCl_2(aq) \ + \ 2AgNO_3(aq) \ \rightarrow \ Mg(NO_3)_2(aq) \ + \ 2AgCl(s)$$

A solution containing 0.001 mol of magnesium chloride reacts with excess silver nitrate. What is the mass of the precipitate formed?

[Molar mass/g mol $^{-1}$: AgCl = 143.4]

- ☑ A 0.072 g
- **B** 0.143 g
- ☑ C 0.287 g
- ☑ D 0.574 g

(Total for Question 5 = 1 mark)

6 When 0.127 g of copper is added to excess silver nitrate solution, the following reaction occurs.

$$Cu(s) + 2AgNO_3(aq) \rightarrow Cu(NO_3)_2(aq) + 2Ag(s)$$

What mass of silver is formed?

[Molar masses/g mol⁻¹: Cu = 63.5 Ag = 107.9]

- **△ A** 0.216g
- **B** 0.254 g
- **☑ C** 0.432 g
- **■ D** 0.863 g

(Total for Question 6 = 1 mark)

DO NOT WRITE IN THIS AREA

NOT WRITE IN THIS AREA

www.chemistryhk.com

7 The molecular formula of phosphorus(V) oxide is P_4O_{10} . What is the percentage by mass of phosphorus in this oxide?

[Molar masses/g mol⁻¹: O = 16.0 P = 31.0]

- A 28.6%
- B 42.9%
- ☑ D 56.3%

(Total for Question 7 = 1 mark)

8 Aluminium reacts with hydrochloric acid.

$$2Al(s) + 6HCl(aq) \rightarrow 2AlCl_3(aq) + 3H_2(g)$$

What is the maximum volume of hydrogen at room temperature and pressure (r.t.p.) that can be formed from 0.135 g of aluminium?

[Molar volume of gas at r.t.p. $= 24\,000\,\mathrm{cm^3}\,\mathrm{mol^{-1}}$ Molar mass Al $= 27.0\,\mathrm{g}\,\mathrm{mol^{-1}}$]

- \triangle **A** 60 cm³
- **■ B** 80 cm³
- \square **C** 120 cm³
- \square **D** 180 cm³

(Total for Question 8 = 1 mark)

9 150 cm³ of ethane is mixed with 700 cm³ of oxygen. The equation for the reaction is

$$C_2H_6(g) + 3\frac{1}{2}O_2(g) \rightarrow 2CO_2(g) + 3H_2O(I)$$

What is the total volume of gas when the reaction is complete? All gas volumes are measured at the same temperature and pressure.

- \square **B** 300 cm³
- \square **D** 475 cm³

(Total for Question 9 = 1 mark)


DO NOT WRITE IN THIS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

www.chemistryhk.com

10 In an experiment to determine the enthalpy change for the reaction between zinc and copper(II) sulfate, a cooling curve was used to estimate the temperature change. The zinc was added to the copper(II) sulfate solution at 3½ minutes and the results were plotted on a graph.



What is the temperature change?

- B 48°C
- ☑ D 68°C

(Total for Question 10 = 1 mark)

do not write in this area

www.chemistryhk.com

11 The enthalpy changes of two reactions are

$$2Fe(s) + 1\frac{1}{2}O_2(g) \rightarrow Fe_2O_3(s) \Delta H^{\ominus} = -824 \text{ kJ mol}^{-1}$$

C(s) +
$$\frac{1}{2}O_2(g) \rightarrow CO(g)$$
 $\Delta H^{\ominus} = -110 \text{ kJ mol}^{-1}$

For the reaction

$$Fe_2O_3(s) \ + \ 3C(s) \ \rightarrow \ 2Fe(s) \ + \ 3CO(g)$$

the enthalpy change is

- B +494 kJ mol⁻¹

(Total for Question 11 = 1 mark)

12 Which change would have a **negative** ΔH value?

- \square **A** $Cl(g) + e^- \rightarrow Cl^-(g)$
- \square **B** $Cl_2(g) \rightarrow 2Cl(g)$
- \square C Na(s) \rightarrow Na(l)
- \square **D** Na(g) \rightarrow Na⁺(g) + e⁻

(Total for Question 12 = 1 mark)

DO NOT WRITE IN THIS

DO NOT WRITE IN THIS AREA

www.chemistryhk.com

13 The combustion of magnesium may be represented by two equations.

$$Mg(s) + \frac{1}{2}O_2(g) \rightarrow MgO(s)$$

$$2Mg(s) + O2(g)$$

$$\rightarrow$$
 2MgO(s)

The units of ΔH for equation (1) are kJ mol⁻¹. The units of ΔH for equation (2) are

- \boxtimes **A** kJ mol⁻¹
- \square **B** (kJ mol⁻¹) ÷ 2
- \boxtimes **C** (kJ mol⁻¹) × 2
- \square **D** $(kJ mol^{-1})^2$

(Total for Question 13 = 1 mark)

14 Which of the species, Ne, F⁻ and Na⁺, have the electronic structure 1s² 2s² 2p⁶?

- A Ne only
- **■ B** Ne and F⁻ only
- ☑ C Ne and Na⁺ only
- Ne, F⁻ and Na⁺


(Total for Question 14 = 1 mark)

O NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

www.chemistryhk.com

15 The graph shows the variation of first ionisation energy with atomic number for successive elements in the Periodic Table.



The elements X, Y and Z are

- **A** alkali metals.
- **B** alkaline earth metals.
- **C** halogens.
- **D** noble gases.

(Total for Question 15 = 1 mark)

16 The electrostatic interactions involved in a covalent bond are electron-electron, nucleus-nucleus and electron-nucleus. What types of interaction occur?

	electron-electron	nucleus-nucleus	electron-nucleus
Α	attraction	attraction	repulsion
В	repulsion	repulsion	attraction
C	attraction	repulsion	attraction
D	repulsion	attraction	attraction

(Total for Question 16 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.

X

X

X

X

DO NOT WRITE IN THIS

DO NOT WRITE IN THIS AREA

www.chemistryhk.com

17 What is the name of the organic compound with the structure shown?

- A 2-chloro-1,1-dimethylpropane
- ☑ B 2-chloro-3,3-dimethylpropane
- □ 1-chloro-2,3-dimethylbutane

(Total for Question 17 = 1 mark)

- 18 Methane is considered a better fossil fuel than coal because methane

 - **B** is mainly obtained from renewable sources.
 - **C** produces less carbon dioxide per kWh of power generated.
 - **D** has no effect on the ozone layer.

(Total for Question 18 = 1 mark)

19 How many σ bonds are there in the organic compound with the skeletal structure shown?



- **A** 2
- **B** 3

(Total for Question 19 = 1 mark)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

AREA


DO NOT WRITE IN THIS

DO NOT WRITE IN THIS AREA

SECTION B

Answer ALL the questions. Write your answers in the spaces provided.

21 Mass spectrometry is used to determine the relative atomic masses of elements. The diagram shows the main features of a mass spectrometer.



- (a) After vaporisation, the sample passes through three stages before reaching the detector.
 - (i) The first stage is ionisation, which occurs at **R**. Describe fully the ionisation process for the element nickel, Ni, writing an equation to illustrate it.

State symbols are not required.

(2)

(::)	December 41.					-
(11)	Describe the	processes	occurring	at S	and	Ι.

(2)

S

T.....



www.chemistryhk.com

(iii) Explain why the sample needs to be ionise	d.
---	----

(1)

(b) The tallest peak in a mass spectrum (called the base peak) is given a height of 100 and the heights of all the other peaks are given relative to the base peak. A sample of the element nickel is analysed in a mass spectrometer and found to have two significant peaks.

m / e	Relative peak height
58	100
60	39.8

(i) Calculate the relative atomic mass of nickel in this sample. Give your answer to one decimal place.

(2)



DO NOT WRITE IN THIS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

www.chemistryhk.com

(ii)	Use the relative peak heights to calculate the percentage abundance of the
	two isotopes in the sample.

(2)

(iii) The mass spectrum of this sample of nickel had a very small peak at m / e = 29. Identify the species responsible for this peak. Write an equation to show how it is formed.

State symbols are not required.

(2)

(c) Mass spectrometry is also used to identify chemical compounds. State **one** application for this use of the technique.

(1)

(Total for Question 21 = 12 marks)




www.chemistryhk.com

- **22** Enthalpy changes of combustion are very important in thermochemistry because they can be used to determine enthalpy changes that cannot be measured directly.
 - (a) Define standard enthalpy change of combustion.

(2)

(b) A class of students used the apparatus below to determine the enthalpy change of combustion for some alcohols.





One student obtained the following results for ethanol.

Measurement	Value
Mass of water/g	250.00
Mass of ethanol used/g	0.55
Temperature rise/°C	9.5

DO NOT WRITE IN THIS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

www.chemistryhk.com

(i) Calculate the energy transferred in the student's experiment.

(1)

Use the expression

Energy transferred (J) = mass of water \times 4.18 \times temperature change

(ii) Calculate the enthalpy change of combustion of ethanol. Give a sign and units with your answer.

(3)

- (c) Most of the students obtained similar results for the enthalpy change of combustion of ethanol. The class mean was $-840 \, \text{kJ} \, \text{mol}^{-1}$ compared with the Data Book value of $-1367 \, \text{kJ} \, \text{mol}^{-1}$.
 - (i) Calculate the percentage error in the mean value obtained by the class compared to the Data Book value.

(1)



(ii) One student suggested that the difference between the stude Data Book value was due to the uncertainties in measuring the Explain why this suggestion is incorrect.	ne masses and temperatures.
No calculation is required.	DO: NOT WRITE IN THIS AF
(iii) Suggest one factor that could have caused the difference bet	
students' values and the Data Book value. Justify your answer	DO: NOT WRITE IN THIS AREA
	DO NOT WRITE IN THIS ARE

DO NOT WRITE IN THIS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

www.chemistryhk.com

- (d) The standard enthalpy change of formation of propan-1-ol cannot be measured directly.
 - (i) Complete the Hess cycle below, which may be used to calculate the standard enthalpy change of formation of propan-1-ol. Add missing enthalpy changes, arrows and species. Include state symbols.

(3)

$$C_3H_8O(I)$$
 + $4\frac{1}{2}O_2(g)$ $\xrightarrow{\Delta H_c^{\oplus} \{\text{propan-1-ol}\}} 3CO_2(g)$ + $4H_2O(I)$ $\xrightarrow{3 \times \Delta H_c^{\oplus} \{\text{C(s)}\}}$

(ii) Use your completed cycle in (d)(i) and the data in the table, to calculate the standard enthalpy change of formation of propan-1-ol.

(2)

Substance	$\Delta H_{\rm c}^{\ominus}$ / kJ mol ⁻¹
carbon	-394
hydrogen	-286
propan-1-ol	-2021

(Total for Question 22 = 16 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

	uestion is about the bonds that chlorine forms in its chemical compounds.	
a) Ch	nlorine forms a covalent bond in its compound with hydrogen.	
(i)	Give the electronic configuration of chlorine using the s p d notation.	(1)
(ii)	Draw a dot-and-cross diagram of hydrogen chloride, showing outer electro	ons only. (1)
(iii	i) Describe fully the formation of the covalent bond in hydrogen chloride in terms of orbital overlap.	(3)
(iii		(3)

DO NOT WRITE IN THIS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

www.chemistryhk.com

(b) Chlorine forms ionic bonds with metals such as sodium and silver. The lattice energies of ionic compounds provide information about their bonds. The table below shows the experimental and calculated values for the lattice energy of sodium chloride and silver chloride.

Compound	Lattice energy / kJ mol ⁻¹	
Compound	Experimental	Calculated
sodium chloride	-780	-770
silver chloride	-905	-833


(i) Draw a dot-and-cross diagram of sodium chloride, showing outer electrons only.

*(ii) Explain why the experimental and calculated values for the lattice energy of sodium chloride are similar whereas those for silver chloride differ significantly.

(Total for Question 23 = 9 marks)

(3)

24 The diagram summarises some of the processes involved in the production of alkanes from crude oil, and their uses.



(a) Name the processes shown in the diagram.

(4)

A.....

B.....

C.....

D.....

(b) State what happens to the compounds present in crude oil during process **A**. Identify the property of the compounds which allows this process to work.

(2)

(c)	The naphtha fraction comprises alkanes with a minimum of four and a maximum
	of ten carbon atoms. Write an equation for the formation of octane and ethene in
	process B . State symbols are not required.

(2)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

www.chemistryhk.com

(d) (i) Write an equation for the reaction occurring in process C . State symbols are not required.	(1)
(ii) Octane is converted into cyclooctane on a large scale. Explain why cyclooctane is added to petrol.	(2)
(e) Using displayed formulae, write a balanced equation for process D . State symbols are not required.	(2)



(Total for Question 24 = 13 marks)

www.chemistryhk.com

25 Alkanes and alkenes react with halogens.

a) 🛚	The reaction	of methane	with chlorine	is a free	radical	substitution.
------	--------------	------------	---------------	-----------	---------	---------------

$$CH_4 + Cl_2 \rightarrow CH_3Cl + HCl$$


(i) State the essential condition for this reaction.

(1)

(ii) The first stage in the mechanism of this reaction is the formation of the chlorine free radical.

Explain fully what a curly half-arrow represents in this equation.

(2)



(2)





DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

www.chemistryhk.com

	very much greater than the amount formed in the termination stage.	(3)
•••••		
•••••		
-) D		
	omine reacts with propene in normal laboratory conditions.	
(1)	State the type and mechanism of this reaction.	(1)
(ii)	Draw the structure of the product of this reaction.	(1)
(ii)	Draw the structure of the product of this reaction.	(1)
(ii)	Draw the structure of the product of this reaction.	(1)
(ii)	Draw the structure of the product of this reaction.	(1)
(ii)	Draw the structure of the product of this reaction.	(1)
(ii)	Draw the structure of the product of this reaction.	(1)
(ii)	Draw the structure of the product of this reaction.	(1)
(ii)	Draw the structure of the product of this reaction.	(1)
(ii)	Draw the structure of the product of this reaction.	(1)
(ii)	Draw the structure of the product of this reaction. (Total for Question 25 = 10 main	



0 (8) (18) 4.0	helium 2	20.2 Ne neon 10	39.9 Ar argon 18	83.8 Kr	krypton 36	131.3	Xenon	4	[222] R	radon 86	rted	_		_
7	(77)	19.0 F fluorine 9	35.5 CL chlorine 17	79.9 Br	bromine 35	126.9	I sodine	23	[210] At	astatine 85	een repo		175 Lu lutetlum 71	[257]
9	(16)	16.0 O oxygen 8	32.1 Sulfur 16	79.0	selenium 34	127.6	Tellurium	25	[508]	potonium 84	116 have b ticated		Yb ytterblum 70	[254]
S	(15)	14.0 N nitrogen 7	31.0 Phosphorus 15	74.9 As	U	121.8	Sb	15	209.0 Bi	bismuth 83	tomic numbers 112-116 hav but not fully authenticated		T T T T T T T T T T T T T T T T T T T	[326]
4	(14)	12.0 C carbon 6	Si Silicon 14	72.6	germanium 32	118.7	S #	20	207.2 Pb	lead 82	tomic nun but not fu	Ì	167 Er erblum 68	[253]
e	(13)	10.8 B boron 5	Al Al aluminium 13	69.7 Ga		114.8	r Indium	46	Z04.4	thallium 81	Elements with atomic numbers 112-116 have been reported but not fully authenticated		Holmium 67	[254]
			(21)	65.4 Zn	30 24 2	112.4	Cd	48	200.6 Hg	mercury 80	Elem		163 Dy dysprosium 66	[251] [254]
			(11)	63.5	copper 29	107.9	Ags	41	197.0 Au	gold 79	[272] Rg noertgentum		Tb terblum 65	[245]
5			(01)	58.7 Ni	nickel 28	106.4	Pq mnipelled	46	195.1 Pt	platinum 78	Ds damstadtum n		Gd Sadolinium 64	[247]
			(6)	58.9	cobalt 27	102.9	Rh	45	192.2 Ir	iridium 77	[268] Mt methrerium		Eu Eu europium 63	[243]
0,1 0,1	hydrogen 1		(8)	55.8 Fe	iron 26	101.1	Ru	44	190.2 Os	osmium 76	HS hasslum n		Sm Smarium 62	[242]
			(2)	54.9 Mn	manganese 25	[86]	Ę	43	186.2 Re	rhenium 75	Bh bohrlum		Pm promethum 61	[237]
		nass ool	(9)	52.0 Cr	chromium r	62.6	5	42	83.8	tungsten 74	Sg seaborgium		Nd ecodymium p	11
	Key	relative atomic mass atomic symbol name atomic (proton) number	(5)	50.9	E	92.9	E	41	180.9 Ta	E	Db dubnium s	ш	Pr Nd pxaecdymium neodymium 59 60	[231]
		relativ ator	(4)	47.9 Ti	E	91.2	Zrztrconium	40	178.5 Hf	hafnium 72	[261] Rf nutherfordum		Cerium p	232
			(3)	45.0	scandium 21	88.9	E	39	138.9 La*	lanthanum 57	AC*			"
7	(2)	9.0 Be beryllium 4	Mg magneslum 12	1.0	F	97.8	Ę	38	137.3 Ba	56 56	Ra radium		* Lanthanide series	
-	(1)	6.9 Li lithium 3	Na Sodium	39.1 K	potassium 19	85.5	Ę	37	132.9 Cs	caestum 55	[223] Fr franclum 87		· Lantha	