www.chemistryhk.com				
Candidate surname	Other names			
Pearson Edexcel International Advanced Level Centre Number Candidate Number				
Wednesday 16 J	anuary 2019			
Morning (Time: 1 hour 30 minutes)	Paper Reference WCH02/01			
Chemistry Advanced Subsidiary Unit 2: Application of Core Principles of Chemistry				
Candidates must have: Scientific cale	Total Marks			

Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.

Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.
- Questions labelled with an asterisk (*) are ones where the quality of your written communication will be assessed
 - you should take particular care with your spelling, punctuation and grammar, as well as the clarity of expression, on these questions.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ▶



P54342A
©2019 Pearson Education Ltd.



SECTION A

Answer ALL the questions in this section. You should aim to spend no more than 20 minutes on this section. For each question, select one answer from A to D and put a cross in the box ⊠. If you change your mind, put a line through the box ₩ and then mark your new answer with a cross ⋈.

- 1 Which of these molecules is planar?
 - A Ethane, CH₃CH₃
 - ☑ B Ethanoic acid, CH₃COOH
 - ☑ C Methanal, HCHO
 - ☑ D Methanol, CH₃OH

(Total for Question 1 = 1 mark)

- 2 Which molecule contains bond angles of both 90° and 120°?
 - \triangle A SF_6
 - \square **B** PCl₅
 - ☑ C BCl₃
 - D BeCl₂

(Total for Question 2 = 1 mark)

- **3** Which molecule has the most polar **bond**?
 - A CH₃CH₂NH₂
 - ☑ B CH₃CH₂OH
 - C CH₃CH₂Cl
 - ☑ D CH₃CH₂I

(Total for Question 3 = 1 mark)



 \mathbf{A}

 \square B

 \mathbb{X} C

 \square D

4 Which is correct for tetrabromomethane, CBr₄?

	Polarity of C-Br bond	Polarity of CBr₄ molecule
⊠ A	polar	polar
⊠ B	non-polar	non-polar
⊠ C	non-polar	polar
⋈ D	polar	non-polar

(Total for Question 4 = 1 mark)

5 In alkanes, increasing the length and branching of the carbon chain both affect the boiling temperature.

Which of the following combination of effects is correct?

Effect on boiling temperature			
Increasing chain length Increasing branching			
increases	decreases		
decreases	increases		
decreases	decreases		
increases	increases		

(Total for Question 5 = 1 mark)

- **6** What is the correct order of boiling temperatures for the hydrogen halides, from the lowest to highest?
 - A HCl, HBr, HI, HF
 - **■ B** HF, HCl, HBr, HI
 - ☑ C HBr, HCl, HF, HI
 - D HI, HBr, HCl, HF

(Total for Question 6 = 1 mark)



■ D Oxygen

(Total for Question 7 = 1 mark)

8 When solid potassium bromide reacts with concentrated sulfuric acid, which substance does **not** form?

- 🖾 A HBr
- \square **B** Br_2
- \boxtimes **C** SO₂
- ☑ D H₂S

(Total for Question 8 = 1 mark)

9 Which equation does **not** represent a disproportionation reaction?

- \square A Cl₂ + H₂O \rightarrow HClO + HCl
- \square **B** $3Cl_2 + 6KOH \rightarrow KClO_3 + 5KCl + 3H_2O$
- \square **C** 2KClO₃ \rightarrow 2KCl + 3O₂
- \square **D** 4KClO₃ \rightarrow 3KClO₄ + KCl

(Total for Question 9 = 1 mark)

10 How does the addition of a catalyst to a reaction affect the shape of the Maxwell-Boltzmann distribution curve?

- A There is no change.
- B The peak moves to the left and is higher.
- ☑ D The total area under the curve increases.

(Total for Question 10 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.

DO NOT WRITE IN THIS AREA

11 This question is about the equilibrium between sulfur dioxide, oxygen and sulfur trioxide in the gas phase.

$$SO_2(g) + \frac{1}{2}O_2(g) \rightleftharpoons SO_3(g) \quad \Delta H^{\ominus} = -98 \text{ kJ mol}^{-1}$$

(a) What are the effects of **decreasing** the temperature?

(1)

	Effect on rate	Effect on equilibrium yield of SO ₃
⊠ A	increases	decreases
⊠ B	decreases	increases
⊠ C	decreases	decreases
⊠ D	increases	increases

(b) What are the effects of **increasing** the pressure?

(1)

	Effect on rate	Effect on equilibrium yield of SO ₃
⊠ A	increases	decreases
⊠ B	decreases	increases
⊠ C	decreases	decreases
⊠ D	increases	increases

(c) The equation for the reaction can also be written as

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$

The enthalpy change for this equation is

(1)

- B -98 kJ mol⁻¹

(Total for Question 11 = 3 marks)

12 A sample of 50 cm³ of ethanol gas is burned completely in 200 cm³ of oxygen.

$$C_2H_5OH(g) + 3O_2(g) \rightarrow 2CO_2(g) + 3H_2O(g)$$

All volumes are measured at a temperature of 400 K and 1 atm pressure.

What is the **total** volume of gas when the reaction is complete?

- \triangle A 150 cm³
- \boxtimes **B** 200 cm³
- \boxtimes **C** 250 cm³
- \square **D** 300 cm³

(Total for Question 12 = 1 mark)

- **13** Which is a tertiary alcohol?
 - A 2-methylbutan-2-ol
 - **B** 2-methylbutan-1-ol

 - D pentan-3-ol

(Total for Question 13 = 1 mark)

14 Excess ammonia in ethanol reacts with 1-bromobutane at high pressure.

Which products could be formed in this reaction?

- \square **A** $C_4H_9NH_2$ and NH_4Br
- \blacksquare **B** $C_4H_9NH_2$ and C_4H_{10}
- \square **C** C_4H_{10} and HBr
- \square **D** C_4H_8 , NH_4Br and HBr

(Total for Question 14 = 1 mark)



15	This qu	uestion is about mechanisms and reaction types.	
	(a) Th	e reaction between an alkane and a halogen to form a halogenoalkane is	(1)
	⊠ A	electrophilic addition.	
	⊠ B	nucleophilic addition.	
	⋈ C	free radical substitution.	
	⊠ D	nucleophilic substitution.	
	(b) The	e reaction between ammonia and a halogenoalkane is	(1)
	⊠ A	electrophilic addition.	
	⊠ B	nucleophilic addition.	
		free radical substitution.	
	⊠ D	nucleaphilic substitution	

(Total for Question 15 = 2 marks)

16 Which of these substances causes the greatest amount of anthro	pogenic global v	warming?
--	------------------	----------

- A Carbon dioxide
- **B** Methane
- C Dinitrogen monoxide
- **D** Dichlorodifluoromethane

(Total for Question 16 = 1 mark)

- 17 In which pair do **both** substances deplete the ozone layer?
 - A Water vapour and carbon dioxide
 - **B** Dichlorodifluoromethane and nitrogen monoxide
 - C Dichlorodifluoromethane and carbon dioxide
 - D Water vapour and nitrogen monoxide

(Total for Question 17 = 1 mark)

TOTAL FOR SECTION A = 20 MARKS



SECTION B

Answer ALL the questions. Write your answers in the spaces provided.

- **18** This question is about diamond, graphite and other carbon structures.
 - (a) (i) In diamond, each carbon atom is covalently bonded to four others, in a three-dimensional structure.

Draw a diagram showing this arrangement.

(1)

(ii) Explain the shape and bond angle of this arrangement of carbon atoms in diamond.

(3)

Shap	oe	 	 	
_				

Bond angle

Explanation



(b)	Graphite consists of a layer lattice, with strong covalent bonds within the layer	S
	and a weaker force between the layers.	

(i) Draw a diagram to show part of **one** layer of graphite with between 13 and 19 carbon atoms, and give the bond angle.

(2)

Diagram

Bond angle

(ii) Name the force between the layers.

(1)

(iii) Give a reason why graphite conducts electricity whereas diamond does not.

(1)

ne one other form of pure ca	rbon.		(1)
ne one other form of pure ca	rbon.		(1)
			(=)
		and its physical propert	ies to
9	letting too hot when re-ente Use your knowledge of the s	etting too hot when re-entering the atmosphere	graphite shield was used on the front of early spacecraft to prevent the letting too hot when re-entering the atmosphere. Use your knowledge of the structure of graphite and its physical propert uggest two reasons for this use.

19	This question is about the structure, properties and reactions of the alcohols, ethano and butan-2-ol.	I
	(a) Ethanol mixes with water in all proportions, but butan-2-ol has limited solubility in water.	
	(i) Name all the intermolecular forces present in these alcohols.	(2)
	*(ii) Explain why butan-2-ol has limited solubility in water.	(2)
	(b) Both alcohols react with sodium.	
	(i) Describe what you see when ethanol reacts with sodium.	(2)
	(ii) Write the equation for the reaction of ethanol with sodium. State symbols are not required.	(2)



	th alcohols are separately heated under reflux with acidified tassium dichromate(VI).	
Aft	er refluxing, the organic product is distilled from each mixture.	
(i)	Name the organic product formed from ethanol and give a chemical test, with the result, to show the functional group present in the product.	(2)
	Name	
Test and re	esult	
(ii)	Give the structure of the organic product formed from butan-2-ol.	
(11)	dive the structure of the organic product formed from butain 2 of.	(1)
(iii)	State the ways in which the infrared spectra of butan-2-ol and its oxidation	
	product would differ. Specific wave numbers are not required.	
	specific wave numbers are not required.	(2)
	(Total for Question 19 = 13 mar	·ks)



- **20** This question is about iodine and some of its compounds.
 - (a) Iodine can be obtained from iodine compounds, such as potassium iodide, by reaction with chlorine.
 - (i) Write the **ionic** equation for the formation of iodine, by the addition of chlorine to aqueous potassium iodide.

 State symbols are not required.

(1)

(ii) The colour of the iodine solution formed is red-brown.

Name an organic solvent that can be added to extract iodine from its aqueous solution, and give the colour of the organic layer.

(2)

Name

Colour

(iii) lodine reacts with sodium thiosulfate solution.

$$I_2(aq) + 2Na_2S_2O_3(aq) \rightarrow 2NaI(aq) + Na_2S_4O_6(aq)$$

State which element is oxidised and which is reduced, giving the relevant changes in oxidation number.

(2)

(ii) lodoalkanes are easily distinguished from chloroalkanes and bror using mass spectrometry because they only give a single molecu Suggest why 1-iodopropane only has one molecular ion peak wh 1-chloropropane and 1-bromopropane both have two molecular	lar ion peak. ereas
(iii) The mass spectrum of 1-iodopropane has a peak at $m/e = 43$. Give the structure of the ion responsible for this peak and state h	ow it is formed.
1-iodopropane and 2-iodopropane are separately dissolved in ethanological hot silver nitrate solution is added. A precipitate forms in each reacti (i) State the colour and name of the precipitate.	
Colour	
(ii) Write the ionic equation, with state symbols, for the formation of	this precipitate. (1)



(d) Aqueous sodium hydroxide reacts with 1-iodopropane to form propan-1-ol.

Draw the mechanism for this reaction. Include curly arrows and relevant lone pairs.

(2)

- (e) Hot concentrated potassium hydroxide in ethanol reacts with 1-iodopropane and 2-iodopropane.The same organic compound is formed in each case.
 - (i) State the type of reaction occurring.

(1)

(ii) **Name** the organic product formed in both reactions.

(1)

(Total for Question 20 = 16 marks)

TOTAL FOR SECTION B = 40 MARKS



SECTION C

Answer ALL the questions. Write your answers in the spaces provided.

21 'Hard' water is water that contains significant concentrations of calcium ions or magnesium ions.

Hard water may be produced when rainwater, containing dissolved carbon dioxide, passes through rocks containing calcium carbonate. Calcium hydrogencarbonate is formed in solution.

Hard water may also be formed when rain water passes through minerals containing calcium sulfate or magnesium sulfate.

Hard water causes two problems.

The first is that it forms a precipitate (scum) when mixed with soap.

The second is that, on heating, it will form a deposit of calcium or magnesium carbonate which reduces the efficiency of heating elements and may lead to blockages in boiler pipes.

- (a) A flame test can be used to detect the metal ion present in the precipitate formed by heating hard water.
- *(i) Explain how metal ions produce a colour in a flame test.

 (3)

 (ii) Give the colour of the flame you would expect to see when calcium ions are present.

 (1)

 (iii) Give the reason why magnesium ions do not produce a flame colour.

 (1)



(b) Write an equation to show how calcium hydrogencarbonate is formed when rainwater, containing dissolved carbon dioxide, reacts with calcium carbonate. State symbols are not required.

www.chemistryhk.com

(1)

(c) Suggest why calcium sulfate and magnesium sulfate can form hard water but barium sulfate cannot.

(1)

(d) Group 2 metal carbonates decompose on heating.

$$MCO_3 \rightarrow MO + CO_2$$

*(i) Explain why calcium carbonate requires stronger heating to decompose than magnesium carbonate.

(3)

(ii) When 10.00 g of a Group 2 metal carbonate is completely decomposed by heating, 1.626 dm³ of carbon dioxide forms, at room temperature and pressure (r.t.p.).

Deduce by calculation the metal ion present.

[Molar volume of a gas at r.t.p. = $24.0 \,\mathrm{dm^3 \,mol^{-1}}$]

(3)

(iii) Calcium oxide dissolves in water to form calcium hydroxide (limewater), Ca(OH)₂(aq).

Write the equation for the reaction of carbon dioxide with limewater. Include state symbols.

(1)



(e) The solubility of calcium hydroxide in water is determined by titration of a saturated solution.

An excess of calcium hydroxide is added to about 100 cm³ of distilled water in a conical flask.

The flask is stoppered, shaken and allowed to stand overnight.

10.0 cm³ portions of this saturated solution are titrated with 0.0500 mol dm⁻³ hydrochloric acid.

$$Ca(OH)_2 + 2HCl \rightarrow CaCl_2 + H_2O$$

(i) Suggest a suitable indicator for this titration and give the colour change you would expect to see at the end-point.

(2)

Indicator

Colour change from ______to ____

(ii) The mean titre is 8.90 cm³.

Calculate the concentration of the saturated calcium hydroxide solution, in g dm⁻³.

(4)

(Total for Question 21 = 20 marks)

TOTAL FOR SECTION C = 20 MARKS TOTAL FOR PAPER = 80 MARKS



BLANK PAGE



BLANK PAGE



The Periodic Table of Elements

								_
0 (8)	(18) 4.0 He hetium 2	20.2 Ne	39.9 Ar argon 18	83.8 Kr	krypton 36	131.3 Xe xenon 54	[222] Rn radon 86	ted
7	(17)	19.0 F fluorine 9	35.5 CI chlorine 17	79.9 Br	bromine 35	126.9 I iodine 53	[210] At astatine 85	een repor
9	(16)	16.0 O oxygen 8	32.1 S sulfur 16	. Se	selenium 34	127.6 Te tellurium 52	Po polonium 84	116 have tricated
2	(15)	14.0 N nitrogen	31.0 Phosphorus 15	74.9 As	arsenic 33	Sb antimony 51	209.0 Bi bismuth 83	tomic numbers 112-116 hav but not fully authenticated
4	(14)	12.0 C carbon	- 5	72.6 Ge	germanium 32	118.7 Sn tin 50	207.2 Pb tead 82	Elements with atomic numbers 112-116 have been reported but not fully authenticated
ъ	(13)	10.8 B boron 5	27.0 Al aluminium 13		gallium 31	Indium 49	204.4 TI thallium 81	ents with
	'		(12)	65.4 Zn	zinc 30	Cd Cadmium 48	200.6 Hg mercury 80	Elem
			(11)	63.5 Cu	copper 29	Ag silver 47	197.0 Au gold 79	[272] Rg roentgenlum 111
			(10)	58.7 Z i	nickel 28	Pd Palladium 46	195.1 Pt platinum 78	Ds darmstachtum r 110
			6	8.3 8.3	cobalt 27	Rh rhodium 45	192.2 Ir iridium 77	[268] Mt meitnerium of 109
	1.0 Hydrogen		(8)	55.8 Fe	iron 26	Ru ruthenium 44	190.2 Os osmium 76	HS Hassium r 108
			6	54.9 Mn	nanganese 25	[98] Tc	Re Reinhenium 75	[264] Bh bohrium 107
		nass ool	9	52.0 Cr	chromium r	95.9 Mo molybdenum t	183.8 W tungsten 74	Sg seaborg/um 106
	Key	relative atomic mass atomic symbol name atomic (proton) number	(5)		vanadium 23	92.9 Nb niobium	180.9 Ta tantalum 73	[262] Db dubnium s
		relativ ator atomic	€	-	titanium 22	91.2 Zr Zirconium 40	178.5 Hf hafnium 72	[261] Rf nutherfordium 104
			(3)	Sc 5.0	scandium 21	88.9 Y yttrium 39	La*	(227) AC* actinium 89
2	(2)	9.0 Be beryllium 4	Mg magnesium 12	1	calcium 20	87.6 Sr strontium	137.3 Ba barium t	[226] Ra radium 88
-	(1)	6.9 Li lithium	0 5 5	39.1 K	potassium 19	85.5 Rb rubidium 37	Cs Caesium 55	[223] Fr Fr francium 87

Lanthanide series

175 Lu lutetium

173 Yb ytterbium

Tm thullium 69

167 **Er** erbium 68

165 **Ho** holmium 67

163 **Dy** dysprosium

159 **Tb** terbium 65

157 **Gd** gadolinium

[147]
Pm
promethium

Pr Pr accodymium

Ge 140

8

[231]

232

144 Nd neodymium

99

4

No nobelium 102

Md mendelevlum n. 101

fermium

Es einsteinium

Cf

[245] BK berkelium 97

(247) Cm outum 96

americium

plutonium

Np neptunium

uranium

Thorium

5

8

6

^{*} Actinide series